

# Therapeutic Drug Monitoring TDM & Managing Narrow Therapeutic Index NTI Drugs — Clinical Treatise 2012

Por Cristiano Ricardo · Publicado em 22/08/2012

University clinical pharmacology treatise by Prof. Cristiano Ricardo on therapeutic drug monitoring (tdm) & managing narrow therapeutic index (nti) drugs (Clinical Pharmacokinetics & Precision Pharmacotherapy), featuring mathematical PK/PD modeling, clinical cases, and colorful schematics.

Versão online e eventuais atualizações:

<https://cristianorichardo.com/post/therapeutic-drug-monitoring-tdm-managing-narrow-therapeutic-index-nti-drugs-2012-08-22>

---

Clinical Pharmacokinetics & Precision Pharmacotherapy · Clinical Pharmacology & Pharmacotherapy

Narrow therapeutic index drugs require precise serum concentration assays paired with Bayesian dosing algorithms to prevent catastrophic toxicity while ensuring clinical efficacy.

## Pharmacological Mechanism & Kinetic Schema

TOXIC ZONE (Above MTC)

THERAPEUTIC WINDOW (MEC to MTC)

SUB-THERAPEUTIC ZONE (Below MEC)

Steady State reached (4-5 half-lives)

## The Steady-State Equilibrium Principle

Upon repeated dosing at constant intervals ( $\tau$ ), drug accumulation occurs until the rate of elimination matches the rate of administration. True steady-state is achieved strictly after 4 to 5 elimination half-lives ( $t_{1/2}$ ). Serum blood draws executed before steady-state attainment yield dangerously misleading pharmacokinetic interpretations.

## Vancomycin AUC/MIC Guided Monitoring

Consensus clinical guidelines have shifted from isolated trough monitoring (15–20 mg/L) to 24-hour area-under-the-curve over minimum inhibitory concentration (AUC<sub>24</sub>/MIC) targets of 400 to 600 mg·h/L. This approach maximizes bactericidal killing against MRSA while

dramatically reducing the incidence of acute kidney injury (AKI).

### **Aminoglycoside Extended-Interval (Once-Daily) Dosing**

Exploiting concentration-dependent bactericidal killing ( $C_{max}/MIC \geq 8-10$ ) and the post-antibiotic effect (PAE), consolidated single-dose regimens achieve high therapeutic bactericidal spikes while allowing extended washout periods where trough levels drop below 1 mg/L, preventing saturable endocytic uptake into renal proximal tubular cells.

Prof. Cristiano Ricardo

Clinical Pharmacologist & Biochemist · Specialist in Pharmacokinetics & Hospital  
Pharmacotherapy · Published on August 22, 2012 at 01:35